

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PECK in these groups it is interesting to note that he is credited with 16 species in *Russula*, 11 in *Pholiota*, and 62 in *Cortinarius*, and this has to do only with Michigan species.—J. M. C.

Seedling anatomy.—Holden and Bexon<sup>25</sup> have begun a series of studies on the anatomy of teratological seedlings. The first paper deals with seedlings of *Cheiranthus Cheiri*, which showed "cotyledonary abnormality ranging from hemitricotyly to tetracotyly." The conclusion was reached that there are at least two methods of cotyledonary increase, cotyledonary fission and dichotomy of the growing point of the cotyledon. A third method is somewhat doubtfully suggested, namely "the downward displacement of one or more epicotyledonary leaves."—J. M. C.

Apogamy in Camptosorus.—Mrs. Brown<sup>26</sup> has described a case of apogamy in *C. rhizophyllus* that occurred in cultures to determine if apogamy could be induced by the modification of external conditions. The apogamous outgrowth was in general a cylindrical process, with some interesting details as to shape and structure, in which a cluster of tracheids appeared. Previous experimental work had indicated that bright light and relative dryness were the factors involved; but in this case low nutrition seemed to be more important than either.—I. M. C.

Tropical species of Eupatorium.—ROBINSON<sup>27</sup> has published the results of a study of *Eupatorium* as displayed in the American tropics. The wealth of species illustrates how much of the flora of the world remains to be discovered. There are 39 new species described, in addition to new varieties. He has included also a revision of the Colombian species, recognizing 93 species distributed among 7 sections. "Keyed recensions" are given also of the species of Venezuela (35) and of Ecuador (50).—J. M. C.

The orchids of Java.—SMITH,<sup>28</sup> in a fifth paper on the orchids of Java, continues to bring to light the remarkably rich orchid flora of that island. He discusses 61 species representing 27 genera, including 38 new species and 2 new genera (*Chroniochilus* and *Saccolabiopsis*).—J. M. C.

A new genus of Compositae.—Pritzel<sup>29</sup> has published a new genus (*Basedowia*) of Compositae from Australia. It resembles *Helichrysum*, as the name (*B. helichrysoides*) suggests. The genus is named for Herbert Basedow, state geologist of South Australia.—J. M. C.

<sup>&</sup>lt;sup>25</sup> HOLDEN, H. S., and BEXON, DOROTHY, Observations on the anatomy of teratological seedlings. I. On the anatomy of some polycotylous seedlings of *Cheiranthus Cheiri*. Ann. Botany 32:513-530. *figs.* 17. 1918.

<sup>&</sup>lt;sup>26</sup> Brown, Elizabeth Dorothy Wuist, Apogamy in *Camptosorus rhizophyllus*. Bull. Torr. Bot. Club **46**:27–30. *pl.* 2. 1919.

<sup>&</sup>lt;sup>27</sup> Robinson, B. L., Contrib. Gray Herb. Proc. Amer. Acad. **54**:235–367. 1918.

 $<sup>^{28}\,\</sup>mathrm{Smith},$  J. J., Die Orchideen von Java. Bull. Jard. Bot. Buitenzorg II. no. 26. pp. 135. 1918.

<sup>&</sup>lt;sup>29</sup> PRITZEL, E., *Basedowia*, eine neue Gattung der Compositen aus Zentral-Australien. Ber. Deutsch. Bot. Gesell. **36**:332-337. pl. 12. 1918.